

Projective representations of k -Galilei group

C.Gonera, P.Kosiński*, P.Maślanka*

Department of Theoretical Physics II

University of Łódź

Pomorska 149/153

90-236 Łódź, POLAND

M.Tarlini

Dipartimento di Fisica, Università di Firenze

and

INFN — Firenze, Italy

Abstract

The projective representations of k -Galilei group G_k are found by contracting the relevant representations of κ -Poincare group. The projective multiplier is found. It is shown that it is not possible to replace the projective representations of G_k by vector representations of some its extension.

I Introduction

There has been some attention paid recently to the deformations of space-time symmetries depending on dimensionful parameter, the so-called κ -symmetries [1]–[11]. They are interesting because they provide rather mild deformation of classical space-time symmetries with dimensionful parameter (cut-off?) naturally built in. It is, of course, still open question whether

*supported by KBN grant 2 P03B 130 12

quantum symmetries provide a proper way of introducing a fundamental energy/length scale into the theory; in particular, special attention should be paid to the problems related to noncocommutativity of coproduct which apparently seems to be in some contradiction with kinematical properties of many-particle systems (see, however, Ref.[9]). In spite of that it could be interesting to study in more detail the properties of κ -deformed space-time symmetries. Some preliminary studies of their physical implications were already undertaken. In particular, Bacry [10] has found that they possess some attractive features from the point of view of general requirements imposed on kinematical symmetries.

In most papers that appeared so far the deformations of Poincare symmetry were studied. However, it seems to be interesting to analyse the deformation of its nonrelativistic counterpart, i.e. the deformed Galilei group. One version of deformed Galilei algebra was studied in Ref.[4] where it was shown to provide the symmetry algebra of one-dimensional Heisenberg ferromagnet. In Ref.[3] another Galilei algebra was found by applying the contraction procedure ($c \rightarrow \infty$, $\kappa \rightarrow 0$, $k \equiv \kappa c$ kept fixed) to κ -Poincare algebra in trigonometric version. The properties of this algebra (in hyperbolic version) as well as the algebra obtained by letting $c \rightarrow \infty$, $\kappa \rightarrow \infty$, $k \equiv \kappa/c$ fixed, were studied in Ref.[8]. Finally, in Ref.[11], the κ -Poincare group was contracted to the k -Galilei group and the latter was shown to be dual to k -Galilei algebra. The bicross-product structure of both was revealed and the projective representations of two-dimensional counterpart of k -Galilei group were constructed.

In the present paper we continue the study of k -Galilei group. In Sec.II the projective multiplier is found by contracting the trivial multiplier on κ -Poincare group. In the $k \rightarrow \infty$ limit it reduces to the standard nontrivial multiplier on classical Galilei group. In Sec.III the unitary projective representations of k -Galilei group are constructed (again by contraction from the representations of κ -Poincare group) and their infinitesimal form is given. The generators of infinitesimal representations form the algebra which is a “central” extension of k -Galilei algebra. The question arises whether this structure can be lifted to the Hopf algebra structure. This question is equivalent to the one whether there exists a “central” extension of k -Galilei group such that the projective representations of the latter are equivalent to the standard (vector) representations of its central extension. In Sec.IV we prove that no such central extension exists. Sec.V is devoted to some conclusions.

Finally, the technicalities are relegated to the Appendices.

We conclude the introduction with short resume of results obtained in Ref.[11].

In order to find the k -Galilei group one can apply the contraction procedure to the κ -Poincare group defined in Ref.[2]. The following convenient parametrization of Lorentz group can be used for contraction procedure (actually, it differs slightly from the one adopted in Ref.[2])

$$\begin{aligned}
\Lambda^0_0 &= \frac{1}{\sqrt{1 - \vec{v}^2/c^2}} \equiv \gamma \\
\Lambda^0_i &= \frac{\gamma}{c} v^i \\
\Lambda^i_0 &= \frac{\gamma}{c} v^i \\
\Lambda^i_j &= (\delta^i_k + (\gamma - 1) \frac{v^i v^k}{\vec{v}^2}) R^k_j \\
RR^T &= I
\end{aligned} \tag{1}$$

as well as

$$a^0 = c\tau$$

The following Hopf algebra G_k (k -Galilei group) is obtained from the contraction $\kappa \rightarrow 0$, $c \rightarrow \infty$, $k \equiv \kappa c$ -fixed:

$$\begin{aligned}
[R^i_j, R^k_l] &= 0, \quad [R^i_j, v^k] = 0, \quad [v^i, v^j] = 0 \\
[a^i, a^j] &= 0, \quad [\tau, a^i] = \frac{i}{k} a^i \\
[\tau, v^i] &= \frac{i}{k} v^i, \quad [\tau, R^i_j] = 0 \\
[v^i, a^j] &= \frac{i}{k} (\frac{1}{2} \vec{v}^2 \delta_{ij} - v^i v^j) \\
[R^i_j, a^k] &= \frac{i}{k} (\delta_{ik} v^m R^m_j - v^i R^k_j) \\
\Delta R^i_j &= R^i_k \otimes R^k_j \\
\Delta v^i &= R^i_j \otimes v^j + v^i \otimes I \\
\Delta a^i &= R^i_j \otimes a^j + v^i \otimes \tau + a^i \otimes I \\
\Delta \tau &= \tau \otimes I + I \otimes \tau \\
(R^i_j)^* &= R^i_j, \quad (v^i)^* = v^i, \quad (a^i)^* = a^i, \quad \tau^* = \tau;
\end{aligned} \tag{2}$$

G_k has a bicross-product structure

$$G_k = T^* \bowtie C(E(3))$$

where $C(E(3))$ is the algebra of functions on classical group $E(3)$ generated by R^i_j and v^i while T^* is defined by

$$\begin{aligned} [\tau, a^i] &= \frac{i}{k} a^i, \quad [a^i, a^j] = 0 \\ \Delta a^i &= a^i \otimes I + I \otimes a^i, \quad \Delta \tau = \tau \otimes I + I \otimes \tau \end{aligned}$$

The k -Galilei algebra $\widetilde{G}_k$, dual to G_k , reads

$$\begin{aligned} [J_i, J_k] &= i\epsilon_{ikl} J_l, \quad [J_i, L_k] = i\epsilon_{ikj} L_j, \quad [J_i, P_k] = i\epsilon_{ikj} P_j \\ [L_i, H] &= iP_i, \quad [L_i, P_j] = \frac{i}{2k} \delta_{ij} \vec{P}^2 - \frac{i}{k} P_i P_j \\ \Delta J_i &= J_i \otimes I + I \otimes J_i \\ \Delta H &= H \otimes I + I \otimes H \\ \Delta L_i &= I \otimes L_i + L_i \otimes e^{-\frac{H}{k}} - \frac{i}{k} \epsilon_{ijk} J_j \otimes P_k \\ \Delta P_i &= I \otimes P_i + P_i \otimes e^{-\frac{H}{k}} \\ P_i^* &= P_i, \quad H^* = H, \quad L_i^* = L_i, \quad J_i^* = J_i \end{aligned} \tag{3}$$

$\widetilde{G}_k$ has also bicrossproduct structure

$$\widetilde{G}_k = T \bowtie U(J, L)$$

where $U(J, L)$ is universal covering of Lie algebra $e(3)$ while T is defined by

$$\begin{aligned} [H, P_i] &= 0, \quad [P_i, P_j] = 0 \\ \Delta H &= H \otimes I + I \otimes H, \quad \Delta P_i = P_i \otimes e^{-\frac{H}{k}} + I \otimes P_i \end{aligned}$$

The duality rules are the same as in classical case.

II Projective multiplier on G_k

In analogy with the classical case one can define projective representation of a quantum group A acting in a Hilbert space H as a map $\rho : H \rightarrow H \otimes A$ satisfying

$$(\rho \otimes I) \circ \rho(\psi) = (I \otimes \omega)((I \otimes \Delta) \circ \rho(\psi)) \tag{4}$$

where ω is a unitary element of $A \otimes A$ (projective multiplier) obeying suitable consistency condition [11].

Two projective representations ρ and ρ' are called equivalent if there exists a unitary element $\zeta \in A$ such that

$$\tilde{\rho} = (I \otimes \zeta)\rho \quad (5)$$

The corresponding multipliers are related by the formula

$$(\zeta \otimes \zeta)\omega = \omega' \Delta(\zeta) \quad (6)$$

Obviously, a multiplier ω is trivial (the representation is equivalent to the vector one) if

$$\omega = (\zeta^{-1} \otimes \zeta^{-1})\Delta(\zeta) \quad (7)$$

In the classical case it is sometimes possible to obtain nontrivial multiplier by contraction [12]. Assume the group $\tilde{G}$ is obtained from G by contraction. Even if G does not admit nontrivial projective multipliers, one can proceed as follows. Let $\zeta(g)$ be a unitary function on G , $\zeta(g)\zeta^*(g) = 1$. Define a trivial multiplier on G

$$\omega(g, g') = \zeta^*(g)\zeta^*(g')\zeta(gg') \quad (8)$$

It can happen that, $\zeta(g)$ being properly chosen, the specific combination of ζ 's appearing on the right hand side of eq.(8) tends to the well defined limit under contraction while $\zeta(g)$ itself has no such a limit. We can then expect that the limiting $\omega(g, g')$ is a nontrivial multiplier on $\tilde{G}$. This is, for example, the case for G being Poincare group and

$$\zeta(\{\Lambda, a\}) = e^{-imca^0} \quad (9)$$

The corresponding multiplier $\omega(g, g')$, eq.(8), gives in the contraction limit $c \rightarrow \infty$

$$\tilde{\omega} = e^{-im(\frac{v^2}{2}\tau' + v^k R^k{}_i a'^i)} \quad , \quad (10)$$

the standard multiplier on Galilei group.

Following the classical case we define the trivial projective multiplier on κ -Poincare group:

$$\omega = (e^{imca^0} \otimes e^{imca^0}) e^{-imc(\Lambda^0_{\mu} \otimes a^{\mu} + a^0 \otimes I)} \quad (11)$$

Our aim is to find the limiting form of ω for $\kappa \rightarrow 0$, $c \rightarrow \infty$, $k \equiv \kappa c$; the κ, c -dependence of m is yet unknown and must be determined from condition that the nontrivial limit exists. To this end we rewrite first ω in a more convenient form making explicit the cancellation of divergent terms. The long and rather tedious calculations reported in Appendix lead to the following expression for ω

$$\omega = e^{i(mc - \kappa \ln(ch(\frac{mc}{\kappa}) + \Lambda^0_0 sh(\frac{mc}{\kappa}))) \otimes a^0} e^{-i\kappa \frac{sh(\frac{mc}{\kappa}) \Lambda^0_k}{ch(\frac{mc}{\kappa}) + \Lambda^0_0 sh(\frac{mc}{\kappa})} \otimes a^k} \quad (12)$$

In order to calculate the limiting value of ω we use the parametrization (1). It is easy to check that in order to obtain the nontrivial limit one can choose the following form of m :

$$m = \frac{-k}{2c^2} \ln(1 - \frac{2Mc^2}{k}) \quad (13)$$

where M is some fixed mass parameter (which we assume to be positive); let us note that for k negative, $m > 0$ and $m \rightarrow 0$ for $c \rightarrow \infty$ while $mc^2\tau$ diverges. Taking the $c \rightarrow \infty$ limit in eq.(12) one obtains

$$\tilde{\omega} = e^{-ik \ln(1 + \frac{M\tilde{v}^2}{2k}) \otimes \tau} e^{-\frac{Mv^k R^k_i}{1 + \frac{M\tilde{v}^2}{2k}} \otimes a^i} \quad (14)$$

which can be also written as

$$\tilde{\omega} = e^{-i(\frac{2k}{M\tilde{v}^2} \ln(1 + \frac{M\tilde{v}^2}{2k}) \otimes I) (\frac{M\tilde{v}^2}{2} \otimes \tau + Mv^k R^k_i \otimes a^i)} \quad (15)$$

This expression is a natural generalization of the one obtained in Ref.[11] for two-dimensional case. In the classical limit $k \rightarrow \infty$ it coincides with the standard multiplier (10).

Let us conclude this section by noting one trouble related to the formula (14). In order to keep the Poincare mass m real we had to assume k negative. This, however, implies that $\tilde{\omega}$ is singular somewhere. On the other hand, with k positive, $\tilde{\omega}$ is everywhere regular.

It seems that this trouble cannot be cured in a simple way. The following argument can be given to support this point of view. Using the results contained in Ref.[18] one easily concludes that the general form of irreducible (co)representation of the κ -Poincare group is obtained by replacing the exponentials on the right-hand side of eq.(16) by

$$e^{-i\kappa \ln(\frac{p_0+C}{A}) \otimes a^0} e^{-\frac{i\kappa p_k}{p_0+C} \otimes a^k}$$

where $A = A(m, c, \kappa)$, $C = C(m, c, \kappa)$ are two real functions subject to the condition $C^2 - A^2 = m^2 c^2$ but otherwise arbitrary. So the question arises whether our trouble can be cured by an appropriate choice of A and C such that obtains, in the $c \rightarrow \infty$, $\kappa \rightarrow 0$ limit, the representation given by formula (19) while $m = m(M, c, k)$ lies, for $k > 0$, in the physical region $m > 0$. This seems not to be possible. Let us put again $\vec{p} = \frac{m}{M} \vec{q}$ and consider the second exponential. It is easy to see that, in order to obtain the proper limiting formula, the following condition should be fulfilled

$$\lim_{c \rightarrow \infty} c^2 (1 + \frac{C}{mc}) = \frac{k}{M}$$

However, due to the condition $C^2 - A^2 = m^2 c^2$, $|C| \geq mc$ and the above equation can be satisfied only provided $C = -mc - \Delta$, $\Delta \geq 0$. Then

$$\lim_{c \rightarrow \infty} \frac{c\Delta}{m} = -\frac{k}{M}$$

which is impossible for $k > 0$, $M > 0$ and $m > 0$.

III Contraction of representations

The unitary representations of the κ -Poincare group were constructed in Ref.[13] (see also [14]). This constructions can be summarized as follows. The representation space is the Hilbert space of square integrable (with respect to the standard measure $d^3\vec{p}/2p_0$) functions over the hiperboloid $p^2 = m^2$ taking their values in the vector space carrying the spin s representation of rotation group (s is assumed to be integer, for s halfinteger one should consider quantum ISL(2, $\mathbb{C}$) group [15] which only amounts to small modifications). The

(right) corepresentation reads

$$\rho : f_i(p_\mu) \rightarrow e^{-i\kappa \ln(ch \frac{mc}{\kappa}) + \frac{p_0}{mc} sh(\frac{mc}{\kappa})} \otimes a^0 e^{\frac{-i\kappa sh(\frac{mc}{\kappa}) p_k}{mcch(\frac{mc}{\kappa}) + p_0 sh(\frac{mc}{\kappa})} \otimes a^k} \cdot D_{ij}(R(p \otimes I, I \otimes \Lambda)) f_j(p_\nu \otimes \Lambda^\nu_\mu); \quad (16)$$

here by $D(R(p \otimes I, I \otimes \Lambda))$ we denote the spin s representation of standard Wigner rotation written as an element of tensor product $H \otimes A$.

It follows from eq.(16) that the whole deformation is contained in translation sector; in other words the representation is obtained by integrating the infinitesimal representation given in Majid–Ruegg basis [16],[17]. In the limit $\kappa \rightarrow \infty$ unitary representations of classical Poincare group are recovered.

In order to find the representations of k -Galilei group we apply again the contraction procedure. To this end we put $\kappa = k/c$ and take $m \equiv m(M, k, c)$ as defined by eq.(13). As in the classical case it is necessary to subtract the rest energy by redefining ρ :

$$\tilde{\rho} \equiv (I \otimes e^{-imca^0})\rho \quad (17)$$

Finally, contrary to the classical case, we have to redefine the momenta and the wave functions as follows:

$$\vec{p} = \frac{m}{M} \vec{q}, \quad \frac{m}{M} f_i(\vec{p}) = \tilde{f}_i(\vec{q}) \quad (18)$$

It is now easy to check that the limit $c \rightarrow \infty$ exists and gives the following unitary representation of k -Galilei group:

$$\tilde{\rho}_{nr} : \tilde{f}_i(\vec{q}) \rightarrow e^{-ik \ln(1 + \frac{\vec{q}^2}{2Mk}) \otimes \tau} e^{\frac{-iq_k}{1 + \vec{q}^2/2Mk} \otimes a^k} (I \otimes D_{ij}(R)) \tilde{f}_j(q_k \otimes R^k_i + I \otimes Mv^k R^k_i) \quad (19)$$

acting in the Hilbert space of functions square integrable with respect to invariant measure $d^3\vec{q}$ and taking values in the vector space carrying spin s representation of rotation group.

As a next step let us find the infinitesimal form of representation ρ_{nr} . Let us recall that if

$$\rho : H \ni f \rightarrow f_{(\alpha)} \otimes a_{(\alpha)} \in H \otimes A$$

is the (right) corepresentation of the quantum group A then any element X of the dual Hopf algebra (quantum Lie algebra) is represented by the operator

$$\tilde{X} : H \ni f \rightarrow f_{(\alpha)} \langle a_{(\alpha)}, X \rangle \in H \quad (20)$$

The relevant duality rules can be, as it was mentioned above, adopted from classical theory. A simple calculation then gives

$$\begin{aligned} J_k &= -i\epsilon_{klm} q_l \frac{\partial}{\partial q_m} + s_k \\ L_k &= iM \frac{\partial}{\partial q_k} \\ H &= k \ln(1 + \frac{\vec{q}^2}{2Mk}) \\ P_k &= \frac{q_k}{1 + \frac{\vec{q}^2}{2Mk}} \end{aligned} \quad (21)$$

Let us note that H and P_k are nonsingular only provided $k > 0$.

The operators (21) verify the following commutation rules

$$\begin{aligned} [J_i, J_k] &= i\epsilon_{ikl} J_l, \quad [J_i, P_k] = i\epsilon_{ikl} P_l, \quad [J_i, L_k] = i\epsilon_{ikl} L_l \\ [K_i, H] &= iP_i \\ [K_i, P_j] &= iM\delta_{ij}e^{-\frac{2H}{k}} + \frac{i}{2k}\delta_{ij}\vec{P}^2 - \frac{i}{k}P_iP_j \end{aligned} \quad (22)$$

For $M \rightarrow 0$ this algebra coincides with the algebraic sector of k-Galilei algebra (3).

Finally, let us note the following dispersion relation valid within the representation (21)

$$k(1 - e^{-\frac{H}{k}}) = \frac{\vec{P}^2}{2M} \quad (23)$$

IV The central extension of k-Galilei group

It is well known that in the classical case given a projective representation of a group G one can construct the group G' such that this projective representation of G is equivalent to the vector representation of G'. The natural

question arises whether the analogous construction is possible in quantum case.

Let us assume that there exists a Hopf algebra G'_k such that

- (i) G'_k is obtained from G_k by adding one new unitary element $\zeta : \zeta\zeta^* = \zeta^*\zeta = I$
- (ii) G_k is a Hopf subalgebra of G'_k
- (iii) $\Delta(\zeta) = (\zeta \otimes \zeta)\omega$

where ω is a projective multiplier.

Then, if ρ is a projective representation of G_k determined by ω (cf. eq.(4)),

$$\rho' = (I \otimes \zeta)\rho$$

is a vector representation of G'_k .

In the commutative case (iii) determines G'_k uniquely and consistently. In the quantum case, however, Δ should be a homomorphism which, together with (iii), imposes nontrivial consistency conditions. It has been already shown [8] that G'_k cannot be obtained by straightforward generalization of standard contraction from trivial extension of κ -Poincare. We show below that there is no solution to the problem, at least if the existence of well-defined limit $k \rightarrow \infty$ which reproduces the classical situation is assumed. To this end let us note first that eqs.(2) and (14) imply

$$e^{\alpha(\tau \otimes I + I \otimes \tau)} \tilde{\omega} e^{-\alpha(\tau \otimes I + I \otimes \tau)} = e^{i \left(\frac{2k}{M\bar{v}^2} \ln(1 + \frac{M\bar{v}^2}{2k} e^{\frac{2i\alpha}{k}}) \otimes I \right) \left(\frac{M\bar{v}^2}{2} \otimes \tau + Mv^k R^k_i \otimes a^i \right)} \quad (24)$$

Therefore

$$[\tau \otimes I + I \otimes \tau, \tilde{\omega}] = \frac{2}{k} \tilde{\omega} (M\bar{v}^2 \otimes \tau + Mv^k R^k_i \otimes a^i) \left(\frac{1}{1 + \frac{M\bar{v}^2}{2k}} \otimes I \right) \quad (25)$$

The homomorphism condition

$$[\Delta(\zeta), \Delta(\tau)] = \Delta([\zeta, \tau]) \quad (26)$$

gives

$$\begin{aligned} \Delta(\zeta) \left(\frac{2}{k} \left(\frac{1}{1 + \frac{M\bar{v}^2}{2k}} \otimes I \right) \left(\frac{M\bar{v}^2}{2} \otimes \tau + Mv^k R^k_i \otimes a^i \right) \right) + \\ + (I \otimes \zeta)([\zeta, \tau] \otimes I) \tilde{\omega} + (\zeta \otimes I)(I \otimes [\zeta, \tau]) \tilde{\omega} = \Delta([\zeta, \tau]) \end{aligned} \quad (27)$$

It follows from eq.(27) that the commutator $[\zeta, \tau]$ should be of the form

$$[\zeta, \tau] = \frac{2}{k} \zeta X \quad (28)$$

where X is an element of G_k , the $1/k$ factor is extracted out explicitly and the factor 2 is written for convenience.

The following relation follows immediately from eqs.(27) and (28)

$$\begin{aligned} \Delta(X) = \tilde{\omega}^{-1}(X \otimes I) \tilde{\omega} + \tilde{\omega}^{-1}(I \otimes X) \tilde{\omega} + \left(\frac{1}{1 + \frac{M\vec{v}^2}{2k}} \otimes I \right) \cdot \\ \cdot \left(\frac{M\vec{v}^2}{2} \otimes \tau + Mv^k R^k_i \otimes a^i \right) \end{aligned} \quad (29)$$

Taking the lowest term in $1/k$ expansion we get

$$\frac{M\vec{v}^2}{2} \otimes \tau + Mv^k R^k_i \otimes a^i = \Delta(X) - X \otimes I - I \otimes X \quad (30)$$

which can be viewed as the relation on classical Galilei group. However, eq.(30) does not hold true because the left-hand side is not a coboundary.

V Conclusion

Using the contraction technique we have found the projective multipliers on k -Galilei group and the corresponding projective representations, both in global as well as in infinitesimal form. It appears that we obtain a well-defined and regular structure for $c \rightarrow \infty$ provided the deformation parameter k is taken to be positive. On the other hand, in order to keep the Poincare mass parameter in the allowed region in the course of contraction we should rather assume k to be negative. We do not have a clear understanding of this phenomenon.

In the classical case the projective representations can be always converted into the vector representations of suitably defined extension of the original group. We have seen in Sec.IV that this is not necessarily the case for quantum groups. There exists no suitable extension of G_k which, in the classical limit $k \rightarrow \infty$ reduces to the standard case. This seems to be not a serious obstacle because it concerns some technical rather than fundamental aspect of the theory.

The problem which certainly deserves further study is the multiplication of representations. This is important if we would like to reconcile the non-commutativity of the algebra coproduct with the basic properties of many-particle systems, especially those containing identical particles.

Appendix

We derive here eq.(12). In order to simplify the notation we omit the tensor product symbol $\otimes$ writing instead with prime the factors appearing right to it. Eq.(11) reads then

$$\omega = e^{imca^0} e^{imca'^0} e^{-imc(\Lambda^0_0 a'^0 + \Lambda^0_k a'^k + a^0)} \quad (\text{A.1})$$

Define

$$X(m) \equiv e^{imca^0} e^{-imc(\Lambda^0_0 a'^0 + \Lambda^0_k a'^k + a^0)} \quad (\text{A.2})$$

Then $X(0)=I$ and $X(m)$ obeys the equation

$$\dot{X}(m) = (-ic)(Y_0(m)a'^0 + Y_k(m)a'^k)X(m) \quad (\text{A.3})$$

where

$$Y_\mu(m) = e^{imca^0} \Lambda^0_\mu e^{-imca^0}, \quad Y_\mu(0) = \Lambda^0_\mu \quad (\text{A.4})$$

Let us first calculate $Y_0(m)$. Using κ -Poincare group commutation rules we get

$$\begin{aligned} \dot{Y}_0(m) &= ice^{imca^0} [a^0, \Lambda^0_0] e^{-imca^0} = -\frac{c}{\kappa} e^{imca^0} ((\Lambda^0_0)^2 - 1) e^{-imca^0} = \\ &= -\frac{c}{\kappa} (Y_0^2(m) - 1) \end{aligned} \quad (\text{A.5})$$

As $Y_0(m)$ commute for all m , (A.5) together with the initial condition (A.4) can be solved by separation of variables yielding

$$Y_0(m) = \frac{\Lambda^0_0 ch(\frac{mc}{\kappa}) + sh(\frac{mc}{\kappa})}{\Lambda^0_0 sh(\frac{mc}{\kappa}) + ch(\frac{mc}{\kappa})} \quad (\text{A.6})$$

With $Y_0(m)$ explicitly known one can apply the same procedure to find $Y_k(m)$; the result reads

$$Y_k(m) = \frac{\Lambda_k^0}{ch(\frac{mc}{\kappa}) + \Lambda_0^0 sh(\frac{mc}{\kappa})} \quad (\text{A.7})$$

Now we can go back to the eq.(A.3). It cannot be solved directly because two terms on the right-hand side do not commute. To account for this we pass to “interaction picture” and define

$$X(m) = e^{-ic \int_0^m dm Y_0(m) a'^0} W(m), \quad W(0) = I \quad (\text{A.8})$$

For (A.3) we get

$$\dot{W}(m) = -ic Y_k(m) e^{ic \int_0^m dm Y_0(m) a'^0} a'^k e^{-ic \int_0^m dm Y_0(m) a'^0} W(m) \quad (\text{A.9})$$

On the other hand

$$ic \int_0^m dm Y_0(m) = i\kappa \ln(ch(\frac{mc}{\kappa}) + \Lambda_0^0 sh(\frac{mc}{\kappa})) \quad (\text{A.10})$$

It follows from (A.9) and (A.10) that

$$\dot{W}(m) = \frac{-ic \Lambda_k^0 a'^k}{(ch(\frac{mc}{\kappa}) + \Lambda_0^0 sh(\frac{mc}{\kappa}))^2} W \quad (\text{A.11})$$

Again everything commutes so that (A.11) can be solved

$$W = e^{-ic \int_0^m \frac{dm}{(ch(\frac{mc}{\kappa}) + \Lambda_0^0 sh(\frac{mc}{\kappa}))^2} \Lambda_k^0 a'^k} \quad (\text{A.12})$$

or

$$W = e^{i\kappa \frac{sh(\frac{mc}{\kappa}) \Lambda_k^0 a'^k}{ch(\frac{mc}{\kappa}) + \Lambda_0^0 sh(\frac{mc}{\kappa})}} \quad (\text{A.13})$$

Eq.(11) follows directly from (A.1), (A.2), (A.8) and (A.13).

References

- [1] J.Lukierski, A.Nowicki, H.Ruegg, Phys.Lett. **B293**, 344 (1992)
- [2] S.Zakrzewski, J.Phys. **A27**, 2075 (1994)
- [3] S.Giller, P.Kosiński, M.Majewski, P.Maślanka, J.Kunz, Phys.Lett. **B286**, 57 (1992)
- [4] F.Bonecchi, R.Giachetti, E.Sorace, M.Tarlini, “*Heisenberg XXZ Model and Quantum Galilei Group*”, hep-th 9204054
F.Bonecchi, E.Celeghini, R.Giachetti, E.Sorace, M.Tarlini, “*Quantum Galilei Group as Symmetry of Magnons*”, hep-th 9203048
F.Bonecchi, E.Celeghini, R.Giachetti, E.Sorace, M.Tarlini, “*Induced representations of the one dimensional quantum Galilei group*”, q-alg 9610006
- [5] J.Lukierski, H.Ruegg, W.J.Zakrzewski, Ann.Phys. **243**, 90 (1995)
- [6] J.Lukierski, H.Ruegg, Phys.Lett. **B329**, 189 (1994)
- [7] P.Maślanka J.Phys. **A 26**, L1251 (1993)
- [8] J.A.De Azcarraga, J.C.Perez Bueno, J.Math.Phys. **36**, 6879 (1995)
J.A.De Azcarraga, J.C.Perez Bueno, J.Phys. **A29**, 6353 (1996)
- [9] P.Kosiński, P.Maślanka, “*The toy model of nonrelativistic quantum space-time symmetry*”, talk given on IV International Wigner Symposium, Wien 1997
- [10] H.Bacry, Phys.Lett. **B306**, 41 (1993)
H.Bacry, J.Phys **A26**, 5413 (1993)
- [11] S.Giller, C.Gonera, P.Kosiński, P.Maślanka, Mod.Phys.Lett. **A10**, 2757 (1995)
- [12] E.J.Saletan, Journ.Math.Phys. **2**, 1 (1961)
- [13] P.Maślanka, Journ.Math.Phys. **35**, 5047 (1994)

- [14] P.Kosiński, J.Lukierski, P.Maślanka, A.Sitarz, “*From differential calculus to relativistic fields on noncommutative κ -deformed Minkowski space*”, in preparation
- [15] P.Maślanka, Journ.Phys. A **A27** , 7099 (1994)
- [16] S.Majid, H.Ruegg, Phys.Lett. **B334**, 348 (1994)
- [17] K.Przanowski, “*Infinitesimal induced representations of κ -Poincare group*”, Proc.IX Max Born Symposium, Karpacz 1996.
- [18] S.Giller, C.Gonera, M.Majewski, Acta Phys. Polon **B27**, 2131 (1996).